

Identification of preoperative and intraoperative predictors of regional cerebral oxygen desaturation during adult elective cardiac surgery

Ioanna Apostolidou, MD¹, Nancy Nussmeier, MD², Muhammad Sarwar, MD², Derek Weiland, MD¹, Mojca Konia, MD¹

¹Anesthesiology, University of Minnesota, Minneapolis, MN, ²Anesthesiology, SUNY Upstate Medical Center, Syracuse, NY.

Introduction

Previous studies indicated that regional cerebral oxygen desaturation is associated with worse major organ morbidity and mortality, postoperative cognitive dysfunction and increased hospital LOS in cardiac surgery (1-3). In a recent study, preoperative rSO₂ was directly related to body mass index, gender, and preoperative hematocrit (Hct), and was inversely related to age, creatinine, LVEF < 30%, and EuroSCORE (4).

Objective

To evaluate associations of preoperative characteristics and intraoperative factors to cerebral oxygen desaturation in adult cardiac surgery patients

Study Design and Subjects

- Prospective observational multicenter study in three clinical sites in the US: University of Minnesota, SUNY Upstate Medical Center, Minneapolis Heart Institute
- 90 adult patients undergoing elective cardiac surgery, aged ≥ 18 years old

Measurements


Preoperative characteristics: age, height, weight, BMI, gender, race, smoking status, prior surgeries, disease vessels, cerebrovascular disease, prior carotid surgery, dialysis and Hct.
 Intraoperative factors included lowest Hct, duration of surgery, duration and use of CPB.

- Univariate and multivariate linear and logistic regression analysis

rSO₂ Measurements

- Cerebral oximetry monitoring was conducted using the Nonin Model 7600 NIRS regional oximetry with 8000CA sensor
- Based on Equanox Regional Oximetry System
- Three wavelengths to improve accuracy
- Dual-emitter to eliminate surface effect
- Critical rSO₂ thresholds used:


- 1) 25% below baseline
- 2) Below 50% absolute rSO₂
- 3) Nadir rSO₂



Results

Baseline characteristics	All sites (n=90)
Age (years)	61.6 ± 12.9
% Male	59/90 (65.6%)
Smoker	14/89 (15.7%)
Height (cm)	172.1 ± 10.3
Weight (kg)	91.3 ± 21.4
BMI	30.1 ± 7.2
Diabetes	33/90 (36.7%)
Redo	14/90 (25.6%)
On Pump	69/86 (80.2%)
Prior CVA	5/90 (5.6%)
Pre-op Dialysis	2/90 (2.2%)

Changes in rSO₂ from Baseline by Timepoint




- 11% of subjects dropped ≥ 25% below the baseline
- 16.7% dropped below 50% rSO₂
- No associations with a 25% drop from baseline

Univariate predictors of dropping below 50% rSO₂

variable	OR	P-value	
Male	4.5	0.01	5/55 M, 9/29 F
Height (cm)	0.898 (1.1)*	0.002	1 cm increase
Weight (kg)	0.954 (1.05)*	0.009	1 kg increase
Preop Hct	0.860 (1.16)*	0.01	1 mg/dL increase
Lowest Hct	0.811 (1.23)*	0.04	1 mg/dL increase

* Denotes the ORs reported in the abstract which were expressed in reverse
Height was a significant predictor in the multivariate model



Univariate predictors of nadir rSO₂

variable	OR	P-value	
Age	-0.1288	0.02	1 yr increase
Height (cm)	0.2118	0.002	1 cm increase
Weight (kg)	0.0985	0.004	1 kg increase
Preop Hct	0.5802	<0.0001	1 mg/dL increase
Lowest Hct	0.4175	0.01	1 mg/dL increase
CPB duration	-0.2869	0.02	Each 10 min


Nadir rSO₂ associations:

Nadir rSO₂ was inversely related (univariate) to:

- age, CPB duration

Nadir rSO₂ was directly related to:

- height, weight, pre-op hematocrit, lowest hematocrit, male gender and smoking
- Age (-0.12/year, p=0.01) and height (0.17/cm p=0.04) remained significant predictors in the multivariate model



Significance

- Factors that determine rSO₂ are cerebral blood flow, hemoglobin, SaO₂, PaCO₂, and cerebral O₂ consumption.
- Our data demonstrated that older and larger patients have paradoxically lower risk of cerebral oxygen desaturation
- Anemia is associated with lower nadir rSO₂ values
- This study supports the obesity paradox.
- This study will help us gain further insight into the nature of patient factors that might affect the variation in cerebral oxygen saturation, and patterns of desaturation.
- This will help us to trigger appropriate preventive measures aiming to improve perioperative cardiac outcome.

References

1. Murkin JM. Anesth Analg 2007; 104: 51-8.
2. Murkin JM. Anesth Analg 2007; 105:536-8.
3. Slater JP. Ann Thorac Surg 2009; 87: 36-44.
4. Heringlake M. Anesthesiology. 2011 Jan;114(1):58-69.

Disclosures

This study has been funded by Nonin Medical, Inc. Plymouth, Minnesota